



Mind-Body Therapies in Children and Youth

SECTION ON INTEGRATIVE MEDICINE

Mind-body therapies are popular and are ranked among the top 10 complementary and integrative medicine practices reportedly used by adults and children in the 2007–2012 National Health Interview Survey. A growing body of evidence supports the effectiveness and safety of mind-body therapies in pediatrics. This clinical report outlines popular mind-body therapies for children and youth and examines the best-available evidence for a variety of mind-body therapies and practices, including biofeedback, clinical hypnosis, guided imagery, meditation, and yoga. The report is intended to help health care professionals guide their patients to nonpharmacologic approaches to improve concentration, help decrease pain, control discomfort, or ease anxiety.

INTRODUCTION

Mind-body therapies and practices (eg, meditation and yoga) are among the top 10 complementary therapies reportedly used by adults and children in the 2007–2012 National Health Interview Survey.¹ Mind-body therapies focus on the interaction between the mind and the body, with the intent to use the mind to influence physical functions and directly affect health. Complementary therapies, such as yoga, meditation, mindfulness-based stress reduction (MBSR), hypnotherapy, guided imagery, and biofeedback, embrace this concept. Data from the 2012 National Health Interview Survey show that 3.7% of US children 4 to 17 years of age used mind-body approaches. Mind-body therapies were used slightly more in older youth aged 13 to 17 years, more than twice as often among females versus males (5.7% vs 1.7%), and less often in the South (2.4%). Children and youth were more likely to use mind-body therapies if they experienced pain-related conditions or emotional, behavioral, or mental conditions and if they received specialty or mental health care. The most common reasons for the use of mind-body approaches were to improve overall health and feel better, to reduce stress level or relax, for general wellness or disease prevention, and to feel better emotionally.² Children are very capable of engaging in self-care skills such as

abstract

FREE

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

Clinical reports from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, clinical reports from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

The guidance in this report does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All clinical reports from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

DOI: 10.1542/peds.2016-1896

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they do not have a financial relationship relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

To cite: AAP SECTION ON INTEGRATIVE MEDICINE. Mind-Body Therapies in Children and Youth. *Pediatrics*. 2016;138(3):e20161896

mind-body therapies,³ and there are many mind-body skills that children and adolescents can learn and apply throughout life.⁴

A growing body of evidence supports the effectiveness and safety of mind-body therapies in pediatrics. In this clinical report, relevant evidence regarding biofeedback, clinical hypnosis, guided imagery, meditation/MBSR, and yoga is reviewed so that pediatric health care providers are better prepared to answer parent questions and provide patient-centered, evidence-based care. For each therapy reviewed, recommendations regarding indications and precautions are provided. The level of evidence based on data from published clinical trials and systematic reviews is described (Table 1). Key outcomes discussed in association with mind-body therapies and practices include focused concentration, decreased pain, and reduced anxiety.

SUMMARIES OF RELEVANT EVIDENCE BY TOPIC

Biofeedback

“Every change in the physiological state is accompanied by an appropriate change in the mental-emotional state, conscious or unconscious, and conversely every change in the mental-emotional state is accompanied by a change in the physiological state.”³³

Biofeedback is defined as the use of electronic or electromechanical equipment to measure and then feed-back information about physiologic processes to an individual. These physiologic processes can then be controlled by the individual for therapeutic purposes. Feedback can be provided in auditory, visual, kinesthetic, or multimedia formats and even now in the form of “video games for the body.”³⁴ This makes biofeedback, in its many forms, particularly relevant as an option for today’s tech-savvy youth.

Although direct clinical observation can provide clues to a patient’s physiologic state and level of autonomic nervous system (ANS) arousal, it is primarily subjective and, therefore, unreliable. In addition, many patients (pediatric, adolescent, or adult) may subjectively state that they “feel relaxed,” but objectively they may not be relaxed at all, at least as defined by measurable physiologic phenomena, especially those that reflect the relative balance of sympathetic and parasympathetic nervous system activity. Therefore, biofeedback can be an invaluable tool for the pediatric health care provider to help gauge what topics, thoughts, and other phenomena trigger mind/body arousal in children and adolescents. The benefits for pediatric patients include allowing them to observe the immediate, convincing, objective mind-body interactions, literally seeing that a “change in the mind (thoughts and/or feelings) can immediately lead to a change in the body’s physiological response.”^{4,35} Interested clinicians can be certified in biofeedback by the Biofeedback Certification International Alliance.^{36,37} The Biofeedback Certification International Alliance certifies individuals who meet education and training standards in biofeedback and progressively recertifies those who advance their knowledge through continuing education.

Research over the past 30 years has shown that children and adolescents are good at self-regulation³⁶ and capable of voluntarily modulating physiologic processes, including peripheral temperature, muscle activity, breathing, brain electrical activity, and certain aspects of immune function, such as salivary immunoglobulin A secretion.³⁵ The most common forms of biofeedback that reflect the balance of the ANS include the following: (1) peripheral temperature (measuring the temperature change in hands or

fingers), (2) heart rate variability (measuring the beat-to-beat variation in heart rate patterns over time), (3) electrodermal activity (measuring sweat gland activity), (4) electromyography (measuring muscle activity), (5) EEG (measuring brain wave activity), (6) capnometry (measuring exhaled carbon dioxide), and (7) pneumography (measuring the movements of the chest and stomach associated with breathing).³⁴

Biofeedback technology has evolved to the point that there are now several low-cost, portable products available that allow for training at home and school, thereby supporting greater generalization of the skill into real-life settings. Enhancing an individual’s context awareness in real-life settings by using biomonitoring and providing real-time feedback is an emerging e-health trend.³⁸ Home biofeedback systems with multimedia game formats are available for personal computers as well as for smart phones and tablet devices.

The user-friendly technologies listed in Table 2 can make practice of these self-regulation skills more enjoyable and effective as they measure a variety of physiologic functions, such as heart rate variability, skin conductance, and peripheral temperature. Like other mind-body skills, it is important that pediatric patients use these skills on a regular basis both for prevention and for acute situational relief, as they eventually learn to control and ultimately reset their ANS response patterns and master the mind-body connection.

Conclusions

Research suggests benefits of peripheral forms of biofeedback for children and adolescents, particularly for headache (tension type and migraine), asthma, enuresis, and rehabilitation applications, as well as EEG biofeedback (neurofeedback)

TABLE 1 Evidence Summary by Topic

| Study, year (design) | Sample Size, <i>N</i> | Age, y | Study Goal | Intervention | Outcomes |
|---|-----------------------|-------------|---|---|---|
| Biofeedback | | | | | |
| Knox et al, 2011 (clinical trial) ⁵ | 24 | 9–17 | Examined changes in anxiety and depression | Heart rate variability biofeedback based on a session-by-session protocol | Biofeedback-assisted relaxation training can be useful in decreasing anxiety and depressive symptoms |
| Palermo et al, 2010 (meta-analysis) ⁶ | 1247 (25 studies) | 9–17 | Quantify the effects of psychological therapies for the management of chronic pain in youth | Cognitive-behavioral therapy, relaxation therapy, and biofeedback | Omnibus cognitive-behavioral therapy, relaxation therapy, and biofeedback all produced significant and positive effects on pain reduction |
| Monastra et al, 2005 (review) ⁷ | N/A | 6–19 | Effects of EEG biofeedback on ADHD | EEG biofeedback | EEG biofeedback was determined to be “probably efficacious” for the treatment of ADHD |
| Eccleston et al, 2002 (systematic review) ⁸ | 808 | 6–18 | Efficacy of psychological therapy of children and adolescents with chronic pain | Variety of biofeedback modalities | Treatments examined are effective in reducing the severity and frequency of chronic pain |
| Clinical hypnosis | | | | | |
| Rutten et al, 2013 (systematic review) ⁹ | 108 | 5–18 | Assess efficacy of HT in pediatric patients with FAP and IBS | Gut-directed HT | Therapeutic effects of HT seem superior to standard medical care in children with FAP or IBS |
| Accardi and Milling, 2009 (systematic review) ¹⁰ | 528 | 3–19 | Effectiveness of hypnosis in reducing procedure-related pain | Hypnosis | Hypnosis was more effective than standard medical care or control at relieving pain in children during medical procedures |
| Vlieger et al, 2007 (RCT) ¹¹ | 53 | 8–18 | Effectiveness of hypnosis for FAP and IBS | 6 sessions of 50 min over a 3-mo period of gut-directed HT | Gut-directed HT is highly effective in the treatment of children with longstanding FAP or IBS |
| Richardson et al, 2006 (systematic review) ¹² | 313 | 3–18 | Effectiveness of hypnosis for procedure-related pain and distress in pediatric patients with cancer | Hypnosis | Hypnosis has the potential to reduce procedure-related pain and distress in pediatric patients with cancer |
| Butler et al, 2005 (RCT) ¹³ | 44 | 4–15 | Examine whether hypnotic relaxation could reduce distress for children who undergo VCUG | Hypnosis | Results indicate significant benefits for the hypnosis group |
| Calipel et al, 2005 (RCT) ¹⁴ | 50 | 2–11 | Efficacy of hypnosis on anxiety and perioperative behavioral disorders | Hypnosis | Hypnosis alleviates preoperative anxiety |
| Guided imagery | | | | | |
| Weigensberg et al, 2014 (RCT) ¹⁵ | 35 | 14–17 | Determine the effects of the mind-body modality of IGI in obese Latino adolescents | 12 weekly sessions of a lifestyle education plus IGI program | The IGI group showed significant reductions in leisure sedentary behavior and increases in moderate physical activity |
| van Tilburg et al, 2009 (pilot study) ¹⁶ | 34 | 6–15 | Test a home-based, guided imagery treatment protocol using audio and video recordings | 2-mo guided imagery treatment | Guided imagery treatment plus medical care was superior to standard medical care only for the treatment of abdominal pain |
| Weydert et al, 2006 (RCT) ¹⁷ | 22 | 5–18 | Evaluated the therapeutic effect of guided imagery for children with recurrent abdominal pain | 4 weekly sessions of guided imagery with progressive muscle relaxation | Significantly greater decrease in the number of days with pain |
| Meditation and MBSR | | | | | |
| Britton et al, 2014 (RCT, pilot) ¹⁸ | 101 | 11.7 (mean) | Effects of a nonelective, classroom-based, teacher-implemented, mindfulness meditation intervention on standard clinical measures of mental health and affect | 6-wk program with daily mindfulness meditation practice | Both control and intervention groups decreased significantly on clinical syndrome subscales and affect but did not differ in the extent of their improvements |
| Sibinga et al, 2014 (RCT) ¹⁹ | 43 | 13–21 | Explore the specific effects of MBSR for urban youth | 8 weekly 2-h MBSR sessions and a 3-h retreat | MBSR did not result in statistically significant differences in self-reported survey outcomes of interest but was associated with qualitative outcomes of increased calm, conflict avoidance, self-awareness, and self-regulation for urban youth |

TABLE 1 Continued

| Study, year (design) | Sample Size, <i>N</i> | Age, y | Study Goal | Intervention | Outcomes |
|---|-----------------------|---------------------|--|--|--|
| Sibinga et al, 2013 (RCT) ²⁰ | 41 | 11–14 | Effects of a school-based MBSR program for young urban males | 12-session programs of MBSR | Results provide cautious support that MBSR enhances self-regulatory processes for urban male youth, including improved psychological symptoms and enhanced coping |
| Sibinga et al, 2016 ²¹ (RCT) | 300 | 12 (mean) | Ameliorate the negative effects of stress and trauma among low-income, minority, middle-school public school students | 12-wk program | MBSR students had significantly lower levels of somatization, depression, negative affect, negative coping, rumination, self-hostility, and posttraumatic symptom severity |
| Barnes et al, 2012 (RCT) ²² | 62 | 15–17 | Impact of TM on LVM in African-American youth at increased risk of development of cardiovascular disease | 15-min TM sessions twice/day for 4 mo | TM decreased LVM index in prehypertensive African-American adolescents |
| Wright et al, 2011 (RCT) ¹⁹ | 121 | 14–15 | Impact on ABP in African-American patients at increased risk of development of essential hypertension | BAM each weekday, 10-min sessions for 3 mo | BAM participants showed significant reductions in self-reported hostility and 24-h systolic ABP |
| Flook et al 2010 (RCT) ²³ | 64 | 7–9 | Evaluate school-based program of MAPs | 30-min MAPs, twice/week for 8 wk | Stronger effect of MAPs on children with executive function difficulties |
| Biegel et al 2009 (RCT) ²⁴ | 102 | 14–18 | Assess the effect of the MBSR program for adolescents with heterogeneous diagnoses in an outpatient psychiatric facility | 8 weekly MBSR classes, meeting 2 h/wk | MBSR may be a beneficial adjunct to outpatient mental health treatment of adolescents |
| Barnes et al, 2004 (RCT) ²⁵ | 100 | 15–17 | Determine the impact of stress reduction on blood pressure in adolescents by the TM program | 15-min TM sessions, twice/day for 4 mo | Beneficial impact of the TM program in youth at risk of the development of hypertension |
| Barnes et al, 2003 (RCT) ²⁶ | 45 | 15–18 | Determine the effect of stress reduction via the TM program on school rule infractions in adolescents | 15-min TM sessions, twice/day for 4 mo | TM program conducted in the school setting has a beneficial effect on absenteeism, rule infractions, and suspension rates |
| Yoga | | | | | |
| Hagins et al, 2013 (RCT) ²⁷ | 30 | 10–11 | Effects of yoga on physiologic response to behavioral stressor tasks | 50 min yoga, 3 times/wk for 15 wk | No significant differences in physiologic responses to behavioral stressors between groups |
| Telles et al, 2013 (RCT) ²⁸ | 98 | 8–13 | Effects of yoga on physical fitness, cognitive performance, self-esteem | 45 min yoga, 5 d/wk for 3 mo | Social self-esteem higher in control versus yoga group, whereas general and parental self-esteem improved |
| Khalsa et al, 2012 (RCT) ²⁹ | 121 | 15–19 | Evaluate potential mental health benefits of yoga for adolescents in secondary school | 30–40 min yoga, 2–3 times/wk for 11 wk | Measures of anger, resilience and fatigue/inertia significantly improved |
| Nidhi et al, 2012 (RCT) ³⁰ | 72 | 15–18 | Efficacy of yoga on glucose metabolism and blood lipid values in adolescent girls with PCOS | 60 min yoga, 7 d/wk for 12 wk | Fasting insulin, fasting blood glucose, and insulin resistance were significantly improved |
| White, 2012 (RCT) ³¹ | 155 | 8–11 | Efficacy of yoga to reduce perceived stress, enhance coping abilities, self-esteem, and self-regulation | 60 min yoga, 1 d/wk for 8 wk, as well as 10 min yoga homework 6 d/wk | Self-esteem and self-regulation increased in both groups, whereas the yoga group reported greater appraisal of stress and greater frequency of coping |
| Mendelson et al, 2010 (RCT) ³² | 97 | 9.7 and 10.6 (mean) | Improve adjustment among chronically stressed and disadvantaged youth | 45 min yoga, 4 d/wk for 12 wk | Significant improvement in the RSQ Involuntary Engagement Scale and component subscales for rumination, intrusive thoughts, and emotional arousal |

ABP, ambulatory blood pressure; ADHD, attention-deficit/hyperactivity disorder; BAM, breathing awareness meditation; HT, hypnotherapy; IGI, Interactive Guided Imagery; LVM, left ventricular mass; MAP, mindful awareness practice; N/A, not available; PCOS, polycystic ovary syndrome; RSQ, Responses to Stress Questionnaire; VCUg, voiding cystourethrography.

for attention-deficit/hyperactivity disorder. Positive evidence for other indications (eg, insomnia, chronic pain syndromes, and anxiety disorders) exists but is not

conclusive. Biofeedback applications for the treatment of functional gastrointestinal tract disorders is an area of particular promise.⁵⁴ Biofeedback offers a particularly

attractive form of self-regulation for today's youth, given their interest in and comfort with technology. There are no significant contraindications to the use of biofeedback, and the

TABLE 2 Demonstrated Safety and Efficacy of Biofeedback-Based Treatments in a Variety of Childhood Conditions

| Biofeedback-Based Treatment Technique | Condition | Reference |
|--|---|--|
| sEMG and peripheral temperature | Migraine, muscle tension, headache | Andrasik and Schwartz, 2006 ³⁹ ; Nestoriuc et al, 2008 ⁴⁰ |
| Variety of biofeedback modalities | Chronic pain syndromes | Eccleston et al, 2002 ⁸ ; Palermo et al, 2010 ⁶ |
| sEMG pelvic floor biofeedback; anorectal EMG biofeedback; manometric feedback | Functional disorders of elimination | Culbert and Banez 2007, ⁴¹ 2008 ⁴² ; Palsson et al, 2004 ⁴³ ; Weydert et al, 2003 ⁴⁴ |
| Sophisticated multichannel sEMG biofeedback | Developmental disabilities and neuromuscular challenges | Bolek, 2006 ⁴⁵ ; Brütsch et al, 2011 ⁴⁶ ; Wang and Reid, 2011 ⁴⁷ ; |
| EEG biofeedback (also termed neurofeedback) | Attention-deficit/hyperactivity disorder | Wang and Reid, 2011 ⁴⁷ ; Monastra et al, 2005 ⁷ ; Vernon et al, 2004 ⁴⁸ |
| Heart rate variability biofeedback | Performance anxiety | Knox et al, 2011 ⁵ |
| Bifrontal sEMG biofeedback | Asthma | Lehrer et al, 2002 ⁴⁹ |
| sEMG biofeedback | Various learning disorders | Carter and Russell, 1985 ⁵⁰ ; Hoy et al 2011 ⁵¹ |
| Specific biofeedback training targeting lowered sympathetic nervous system arousal | Sleep disorders | Barowsky et al, 1990 ⁵² ; Morin et al, 2006 ⁵³ |

sEMG, surface electromyography.

only barrier may be financial in that both home and professional health care biofeedback hardware/software packages can be somewhat expensive and third-party health care insurance payers do not consistently cover biofeedback treatment. A selection of resources for pediatric health care providers is provided in Table 3.

Clinical Hypnosis

Clinical hypnosis in children and adolescents has seen a surge in both research and clinical application in the past 30 years, although its use in children dates back >200 years.⁵⁵ Hypnosis is defined variably by several professional societies.⁵⁶ Perhaps best stated, “when we are in hypnosis, we intensify our attention, decrease our peripheral awareness and become more receptive to new ideas and associations whenever we reinforce, rewire, reframe or otherwise alter the neurophysiological networks we call ‘experience.’ Trance is what happens when we engage in changing our minds.... Hypnosis is a skill set involving interpersonal communication designed to facilitate therapeutic change in maladaptive psycho-physiological reflexes.”⁵⁶ Pediatric health care providers should understand that hypnotherapy in children is a well-established therapeutic modality, and it should not be confused with or misperceived as the

inappropriate practice of hypnosis by entertainers.

Clinical hypnosis, when provided by appropriately trained individuals, is an adjunctive therapy that can be used by pediatric health care providers to assist in managing conditions that they are already otherwise licensed to treat. For example, a pediatrician may use clinical hypnosis to help a child dealing with enuresis, irritable bowel syndrome (IBS), or anxiety. A licensed mental health practitioner may use clinical hypnosis to help children with anxiety, depression, or posttraumatic stress disorder (PTSD). However, a mental health practitioner should not use clinical hypnosis for a child with IBS without physician comanagement, and pediatricians should not use this technique for PTSD without collaborating with a mental health practitioner.

Case series and clinical trials of clinical hypnosis first appeared in the 1970s and 1980s. Since then, the literature has grown and includes clinical trials, Cochrane reviews, and neuroimaging studies.⁵⁵⁻⁵⁹ However, high-quality randomized controlled trials (RCTs) with clear methodologies remain lacking. Clinical hypnosis involves establishing a strong rapport with patients and individualizing the therapy to the specific goals and

characteristics of the patient. This situation precludes a standardized approach, and large randomized studies of individualized approaches are difficult to conduct.

Functional Abdominal Pain

A few studies have evaluated the effectiveness of hypnosis for functional abdominal pain (FAP) and IBS. Vlieger et al¹¹ randomly assigned 52 children to either hypnosis with an experienced clinician or standard care, which included dietary guidance, medication as needed, and supportive counseling. Twelve sessions over 3 months led to marked improvement in pain frequency and severity in patients in the hypnosis group compared with control patients at the end of the intervention and at 1-year follow-up.¹¹ A later follow-up study at 5 years showed a significantly higher remission rate in the hypnosis group compared with the control group (68% vs 20%; $P = .005$).⁶⁰ A systematic review of 3 trials for FAP and IBS in children and adolescents showed superior efficacy over standard care.⁹

Pain Management

Numerous small trials have shown the efficacy of clinical hypnosis for procedural as well as for chronic pain. Butler et al¹³ compared hypnosis with breathing and relaxation techniques for procedure-related pain and anxiety during

TABLE 3 Resources by Topic

| Resources | Web Sites, Books, DVDs, etc |
|--|---|
| Biofeedback | |
| Biofeedback Certification International Alliance | www.bcia.org |
| Inner Balance | www.heartmath.com |
| Interactive games (Healing Rhythms, Journey to Wild Divine) | www.wilddivine.com |
| “eSense” temperature and sweat gland activity (electrodermal activity) sensors | http://www.mindfield.de/en/biofeedback/products/esense/esense-skin-response |
| Tinke | http://www.zensorium.com/tinke/ |
| Hypnosis | |
| Video: “magic glove” technique, performed by Dr Leora Kuttner ^a | http://www.youtube.com/watch?v=cyApK8Z_SQ0 |
| Dr Laurence Sugarman’s 70-minute DVD ^a | Sugarman L. <i>Hypnosis in Pediatric Practice: Imaginative Medicine in Action</i> [DVD and booklet]. Carmarthen, United Kingdom: Crown House Publishing; 2006 |
| National Pediatric Hypnosis Training Institute | www.nphti.org |
| American Society of Clinical Hypnosis | www.asch.net |
| Society for Clinical and Experimental Hypnosis | www.sceh.us |
| Guided imagery | |
| Health Journeys: Guided Meditation and Imagery | http://www.healthjourneys.com |
| Kaiser Permanente: Guided Imagery (podcasts) | https://healthy.kaiserpermanente.org/health/care/!ut/p/a0/FchBD0MgEADAt_iAzYZEYfFmhH6hhdsGiZIIgELt99seZ9DjC33h0-3cUy18_uxCLD22md9bqnCnLVZ8okd_Nd4zoysVAocj_o9bT-GM6IzVap2MBamIBCGsgEWPBohoUkKp8UErXjnTZxmGL2IKPpl/ |
| Shambala Kids: Guided Imagery and Relaxation Audio CDs | http://shambalakids.com/index.php?option=com_content&view=category&id=40&Itemid=419&lang=us |
| Stress Free Kids Indigo Dreams Audio CDs | http://www.stressfreekids.com/category/cds/children-cds |
| Academy for Guided Imagery | http://acadgi.com |
| Meditation and MBSR | |
| For health professionals: Mind-Body STREAM program | https://mind-bodyhealth.osu.edu |
| For parents: Everyday Blessings: The Inner Work of Mindful Parenting | Kabat-Zinn M, Kabat-Zinn J. <i>Everyday Blessings: The Inner Work of Mindful Parenting</i> . New York, NY: Hyperion; 1998 |
| Growing Happiness | http://growing-happiness.com/mindfulness-training-for-parents/ |
| Meditation for children | http://www.freemeditation.com/online-meditation/meditation-for-children/ |
| For teens: Mindfulness for Teens | http://mindfulnessforteens.com |
| Mindfulness Retreats | http://ibme.info/ |
| Learning to Breathe | http://learning2breathe.org/ |
| UCSD Center for Mindfulness Professional Training | http://mbpti.org/ |
| UCSD Center for Mindfulness blog | https://ucsdcfm.wordpress.com/tag/ron-epstein/ |
| UMass Medical School Center for Mindfulness | http://www.umassmed.edu/cfm/training/ |
| UCLA Mindful Awareness Research Center | http://marc.ucla.edu/ |
| Yoga | |
| Global Family Yoga | http://globalfamilyyoga.com/ |
| International Association of Yoga Therapists | http://iayt.org/ |
| Kripalu Yoga in the Schools | http://nccam.nih.gov/health/yoga |
| Yoga Alliance | http://www.yogaalliance.org/ |
| Yoga Calm | http://www.yogacalm.org/ |
| Yoga for the Special Child | http://www.specialyoga.com/ |
| Yoga in Schools | http://yogainschools.org |
| YogaKids | http://yogakids.com/ |

STREAM, Skills Training for Resilience, Effectiveness, and Mindfulness; UCLA, University of California, Los Angeles; UCSD, University of California, San Diego; UMass, University of Massachusetts.

^a The technique should never be used for entertainment, should only be used by appropriately trained providers, and should only be used in clinical situations for which the provider already has competence in treating.

voiding cystourethrography. Moderate effect sizes for symptom reduction were noted by parents, medical staff, and research observers; and procedure length was reduced by 14 minutes in the hypnosis group. A trial comparing preprocedure

hypnosis with midazolam for anesthesia for abdominal surgery showed reduced anxiety at the time of induction as well as improved behavior outcomes at 1 and 7 days after surgery for the hypnosis group.¹⁴ Two systematic reviews

concluded that there is promising evidence for hypnosis for acute procedure-related pain.^{10,12}

Conclusions

Research suggests benefits of clinical hypnosis for children and

adolescents, particularly for FAP, IBS, and pain management. Promising evidence for its application for other indications (eg, enuresis, tics/Tourette syndrome, migraine, and anxiety) exists but is not conclusive. There are few absolute contraindications to the use of hypnosis. The technique should be used only by appropriately trained providers and in clinical situations in which the provider already is competent managing without the inclusion of hypnosis. A selection of practical resources (eg, videos) for pediatric health care providers is provided in Table 3.

Guided Imagery

Guided imagery is a powerful mind-body technique that invokes all of the senses (sight, sound, taste, touch, smell, and movement). Imagery has a rich history in healing traditions throughout the world. Guided imagery and clinical hypnosis have significant overlap, and many studies combine these modalities. Strengths of guided imagery treatment include that it is not invasive and has flexibility of use in different age ranges (preschool-aged through adolescents and adults) and in various settings (outpatient, inpatient, and acute care). The use of guided imagery has been shown to produce measurable physiologic changes in stress and immune biomarkers.⁶¹ Challenges in the use of imagery include variable training, acceptance of a novel therapy by patient and practitioner, familiarity with and access to high-quality resources, and relative lack of randomized controlled outcome studies in children. Due caution is indicated in patients with a history of physical, sexual, or emotional abuse or those with PTSD, in which case coordination of care with a qualified mental health expert is strongly advisable.

An evidence base for the use of guided imagery in adults is present,

and an evidence base in children is growing. In 1 adult RCT, guided imagery in 96 patients with newly diagnosed breast cancer showed significant correlation with improved mood and quality of life.⁶² For example, a second adult RCT that used guided imagery was correlated with an increase in numbers and activity of beneficial immune function (T helper cells, natural killer cells, lymphokine-activated killer cells, and favorable interleukin-1 β levels) in 80 patients with breast cancer in active treatment.⁶³ An example of a pediatric RCT that used guided imagery involved a 12-week lifestyle intervention trial in 29 Latino adolescents with obesity, in which weekly interactive guided imagery sessions were associated with a statistically significant reduction in salivary cortisol, improved physical activity, and promotion of health behavior change in the treatment group.¹⁵ A second RCT combined guided imagery with progressive muscle relaxation in 22 children ages 5 to 18 years with a diagnosis of recurrent abdominal pain. Guided imagery with progressive muscle relaxation in 4 weekly sessions was associated with a statistically significant reduction in days with pain throughout the 2-month follow-up period.¹⁷ Home-based audio-recorded guided imagery also has been shown to be effective in the reduction of recurrent abdominal pain in a treatment group of 34 children ages 6 to 15 years who were randomly assigned to receive guided imagery versus standard care. Results were maintained throughout the 6-month follow-up period.¹⁶

In addition to these RCTs, other small studies that showed efficacy of guided imagery have been conducted for a variety of medical conditions, including asthma,⁶⁴ sickle cell disease,⁶⁵ procedural anxiety,⁶⁶ and posttraumatic stress.⁶⁷ Both imagery and hypnosis may be combined successfully with other mind-body

therapies, such as biofeedback, to enhance relaxation.⁶⁸ Regulated training for guided imagery does not exist at this time.

Conclusions

Guided imagery appears to be a promising complementary therapy for children and adolescents, with very low reports of adverse effects. Guided imagery as a therapeutic intervention has been shown to have positive effects on psychological functioning, stress reduction, and pain management. Caution is advised in patients with a history of previous emotional, sexual, or physical abuse to avoid an unintended triggering of posttraumatic stress symptoms. Consultation with a mental health practitioner is advised if questions about appropriateness of use exist in this context. More RCTs in children are needed for this noninvasive therapy in the pediatric clinical setting. A selection of guided imagery Web resources is provided in Table 3.

Meditation and MBSR

Meditation

Meditation practices for children and youth have become increasingly popular in schools and medical settings alike. Meditation is the practice of intentional attention training and consists of a number of different specific approaches. Research on meditation in children and youth consists primarily of 2 types of meditation: mindfulness meditation and concentration meditation.

Research on meditation in diverse populations of adults has accumulated sufficiently to provide convincing high-level evidence for reproducible benefits of meditation in mental health and pain management.⁶⁹⁻⁷¹ In addition, data suggest that greater levels of mindfulness in adulthood may mitigate some of the negative health effects of adverse childhood experiences.⁷² The literature in

children and youth, however, is less developed and, although suggestive of benefit, is just beginning to emerge.⁷³⁻⁷⁶ To provide the highest level of available evidence regarding the specific effect(s) attributable to meditation instruction for children and youth, conclusions in this report are based on findings from RCTs with active control conditions.

Mindfulness Meditation

Mindfulness meditation is aimed at enhancing individuals' innate capacity to be purposefully aware of their present-moment emotional, cognitive, and sensory experiences. Through instruction in formal and informal meditation techniques, this capacity for purposeful, moment-by-moment, nonjudgmental awareness develops, along with the ability to shift attention. Several RCTs in youth have evaluated the MBSR program, which has established instructor training through the University of Massachusetts School of Medicine's Center for Mindfulness and has been well researched in adults.⁷⁷ Youth-adapted MBSR programs have been found to be beneficial in improving mental health symptoms, coping, and self-regulatory processes and decreasing blood pressure when used in both primary prevention^{18-21,78,79} and treatment settings.⁸⁰ In children 7 to 9 years of age, an RCT of school-based mindful awareness practice instruction versus a reading program did not show differences by treatment group overall but did reveal improvements in mindful awareness practice participants in executive function among children with lower executive function skills at baseline.²⁴

Concentration Meditation

Concentration meditation involves focusing attention on 1 specific thing, such as a word, phrase, or object.²³ A Cochrane review found current research inadequate to suggest meditation for attention-deficit/

hyperactivity disorder and suggested additional trials.⁸¹ Active-controlled RCTs and active-control programs of concentration meditation in children and youth have included both transcendental meditation (TM) and the relaxation response. Compared with active-control programs, TM has been shown to lead to decreases in blood pressure and left ventricular hypertrophy among African-American adolescents with prehypertension^{22,25} as well as fewer negative school behaviors, such as absenteeism.²⁶ Relaxation response has been associated with improvements in self-esteem.⁸²

Conclusions

Research on structured meditation programs for children and youth is suggestive of benefits, particularly related to improvements in mental health, coping, and self-regulation as well as decreasing hypertension and negative school behaviors. Although there are structured training and certification programs for a number of meditation programs (including MBSR, TM, and mindfulness-based cognitive therapy), there is no formal credentialing or licensure for meditation instruction. Costs vary depending on the format of instruction and are increasingly, but not universally, covered by insurance. Although these results are encouraging, careful attention should be paid to elements of implementation and dissemination to maintain high-quality, effective meditation instruction for children and youth.

Yoga

The word yoga is derived from the Sanskrit word *yuj* meaning "union." An ancient Indian practice, yoga has been classified by the National Center for Complementary and Integrative Health as a mind-body medicine modality.⁸³ According to the 2007 National Health Interview Survey,⁸⁴

yoga was the fifth most commonly used complementary therapy practice among all children ages 2 to 17 years, with ~1.5 million children practicing yoga in the previous year. In a survey of children and adolescents with chronic pain, yoga was preferred by 32% as their first choice of complementary therapy.⁸⁵ Therapeutic yoga is the practice of uniting the mind, the body, and the spirit through mindfulness of breathing and body postures to improve stress coping, lessen pain, and improve specific health conditions. Although not completely understood, yoga effects changes in the parasympathetic nervous system, positively affecting heart rate variability.⁸⁶

Fourteen controlled studies^{27-32,87-94} and 4 systematic reviews⁹⁵⁻⁹⁸ were identified, and all uncontrolled trials and those in which yoga was not the sole treatment intervention⁹⁹ were eliminated from consideration. The conclusions of the systematic reviews can be summarized as follows: yoga appears to be a promising complementary therapy for children and adolescents, especially for those with pain and emotional, mental, and behavioral conditions, with very few reported adverse effects. However, a lack of methodologic and statistical rigor, including small sample sizes, absence of randomization, and a high degree of variability between intervention methods, limits the ability to recommend yoga as a primary intervention for any particular population. On the basis of the 14 individual controlled studies, yoga appears to be a promising complementary therapy and stress-management tool for children and adolescents, with very low reports of adverse effects. Yoga as a therapeutic intervention has positive effects on psychological functioning, especially in children coping with emotional, mental, and behavioral health problems. Specifically, research has shown that educational

curricula incorporating stress-management programs improve academic performance, self-esteem, classroom behaviors, concentration, and emotional balance,⁹⁸ suggesting that schools may be an ideal setting to bring yoga to a heterogeneous, socioeconomically diverse sample of children. In addition, in 4 controlled trials, yoga was shown to positively influence metabolic and hormonal variables. Given the increasing prevalence of obesity and metabolic dysfunction in children, coupled with the relative safety and cost-effectiveness of yoga as an intervention, more research in this population is needed. Limitations of reviewed studies include small sample sizes, high attrition rates, lack of evaluator blinding, reliance on self-report measures, and heterogeneity of intervention and control designs. Well-designed controlled trials of yoga for conditions with strong stress-modulated components are warranted. Excellent candidate conditions include asthma, IBS, inflammatory bowel diseases, juvenile idiopathic arthritis, and fibromyalgia. Given the preference for yoga in studies in children with chronic pain, coupled with biological plausibility for response, limited potential for adverse effects, and promising pilot data, there is a great need for controlled studies in this population.

Conclusions

Yoga appears to be a promising complementary therapy and stress-management tool for children and adolescents, with very low reports of adverse effects. Yoga, as a therapeutic intervention, has positive effects on psychological functioning, especially in children coping with emotional, mental, and behavioral health problems. Yoga generally is not billed for and reimbursed as an insurance-covered therapy. Yoga instructors and centers establish a fee for service (per session or as a package for a set number of classes) on the basis of

community-established standards. The Yoga Alliance¹⁰⁰ sets guidelines for yoga teacher certification in the United States. A selection of yoga Web resources is provided in Table 3.

CONCLUSIONS AND RECOMMENDATIONS

This report examines the best-available evidence for a variety of mind-body therapies and practices in children and youth, including biofeedback, clinical hypnosis, guided imagery, meditation, and yoga. The evidence varies in terms of quantity and quality but generally is supportive of mind-body therapies and practices as safe and potentially effective in common and debilitating conditions, including pain and anxiety. Additional potential benefits for school-aged children include improved concentration and self-esteem. Pediatric health care providers are encouraged to facilitate an open dialog with their patients about their use of complementary therapies and to become familiar with mind-body therapies and practices as nonpharmacologic options to improve mood, behavior, and quality of life, which are of great interest and relevance to children, youth, and their parents/caregivers.

LEAD AUTHORS

Sunita Vohra, MD, FAAP, Past Chairperson, AAP Section on Integrative Medicine
Hilary McClafferty, MD, FAAP

CONTRIBUTING AUTHORS

David Becker, MD, FAAP
Christina Bethell, PhD, MBA, MPH
Timothy Culbert, MD, FAAP
Susanne King-Jones, PhD
Larry Rosen, MD, FAAP
Erica Sibinga, MD, FAAP

SECTION ON INTEGRATIVE MEDICINE EXECUTIVE COMMITTEE, 2014–2015

Hilary McClafferty, MD, FAAP, Chairperson
Erica Sibinga, MD, FAAP
Michelle Bailey, MD, FAAP
Timothy Culbert, MD, FAAP
Joy Weydert, MD, FAAP
Melanie Brown, MD, FAAP

STAFF

Teri Salus, MPA, CPC

ABBREVIATIONS

ANS: autonomic nervous system
FAP: functional abdominal pain
IBS: irritable bowel syndrome
MBSR: mindfulness-based stress reduction
PTSD: posttraumatic stress disorder
RCT: randomized controlled trial
TM: transcendental meditation

REFERENCES

1. Black LI, Clarke TC, Barnes PM, Stussman BJ, Nahin RL. Use of complementary health approaches among children aged 4-17 years in the United States: National Health Interview Survey, 2007-2012. *Natl Health Stat Rep*. 2015;(78):1–19
2. Data Resource Center for Child and Adolescent Health. The National Health Interview Survey (NHIS). Child Complementary and Alternative Medicine (CAM) supplement. Data Resource Center for Child and Adolescent Health; 2012. Available at: <http://childhealthdata.org/learn/nhis>. Accessed December 10, 2015
3. Sussman D, Culbert T. Pediatric self-regulation. In: Levine MD, Carey WB, Crocker AC, eds. *Developmental-Behavioral Pediatrics*. 3rd ed. Philadelphia, PA: WB Saunders; 1999:911–922
4. Kohen DP. A pediatric perspective on mind-body medicine. In: Culbert T, Olness K, eds. *Integrative Pediatrics*. New York, NY: Oxford University Press; 2009:267–301
5. Knox M, Lentini J, Cummings T, McGrady A, Whearty K, Sancrant L. Game-based biofeedback for paediatric anxiety and depression. *Ment Health Fam Med*. 2011;8(3):195–203
6. Palermo TM, Eccleston C, Lewandowski AS, Williams AC, Morley S. Randomized controlled trials of psychological therapies for management of chronic pain in children and adolescents: an updated meta-analytic review. *Pain*. 2010;148(3):387–397

7. Monastra VJ, Lynn S, Linden M, Lubar JF, Gruzelier J, LaVaque TJ. Electroencephalographic biofeedback in the treatment of attention-deficit/hyperactivity disorder. *Appl Psychophysiol Biofeedback*. 2005;30(2):95–114
8. Eccleston C, Morley S, Williams A, Yorke L, Mastroyannopoulou K. Systematic review of randomised controlled trials of psychological therapy for chronic pain in children and adolescents, with a subset meta-analysis of pain relief. *Pain*. 2002;99(1–2):157–165
9. Rutten JM, Reitsma JB, Vlieger AM, Benninga MA. Gut-directed hypnotherapy for functional abdominal pain or irritable bowel syndrome in children: a systematic review. *Arch Dis Child*. 2013;98(4):252–257
10. Accardi MC, Milling LS. The effectiveness of hypnosis for reducing procedure-related pain in children and adolescents: a comprehensive methodological review. *J Behav Med*. 2009;32(4):328–339
11. Vlieger AM, Menko-Frankenhuis C, Wolfkamp SC, Tromp E, Benninga MA. Hypnotherapy for children with functional abdominal pain or irritable bowel syndrome: a randomized controlled trial. *Gastroenterology*. 2007;133(5):1430–1436
12. Richardson J, Smith JE, McCall G, Pilkington K. Hypnosis for procedure-related pain and distress in pediatric cancer patients: a systematic review of effectiveness and methodology related to hypnosis interventions. *J Pain Symptom Manage*. 2006;31(1):70–84
13. Butler LD, Symons BK, Henderson SL, Shortliffe LD, Spiegel D. Hypnosis reduces distress and duration of an invasive medical procedure for children. *Pediatrics*. 2005;115(1). Available at: www.pediatrics.org/cgi/content/full/115/1/e77
14. Calipel S, Lucas-Polomeni MM, Wodey E, Ecoffey C. Premedication in children: hypnosis versus midazolam. *Paediatr Anaesth*. 2005;15(4):275–281
15. Weigensberg MJ, Lane CJ, Ávila Q, et al. Imagine HEALTH: results from a randomized pilot lifestyle intervention for obese Latino adolescents using Interactive Guided ImagerySM. *BMC Complement Altern Med*. 2014;14:1–13
16. van Tilburg MA, Chitkara DK, Palsson OS, et al. Audio-recorded guided imagery treatment reduces functional abdominal pain in children: a pilot study. *Pediatrics*. 2009;124(5). Available at: www.pediatrics.org/cgi/content/full/124/5/e890
17. Weydert JA, Shapiro DE, Acra SA, Monheim CJ, Chambers AS, Ball TM. Evaluation of guided imagery as treatment for recurrent abdominal pain in children: a randomized controlled trial. *BMC Pediatr*. 2006;6:1–10
18. Britton WB, Lepp NE, Niles HF, Rocha T, Fisher NE, Gold JS. A randomized controlled pilot trial of classroom-based mindfulness meditation compared to an active control condition in sixth-grade children. *J Sch Psychol*. 2014;52(3):263–278
19. Sibinga EM, Perry-Parrish C, Thorpe K, Mika M, Ellen JM. A small mixed-method RCT of mindfulness instruction for urban youth. *Explore (NY)*. 2014;10(3):180–186
20. Sibinga EM, Perry-Parrish C, Chung SE, Johnson SB, Smith M, Ellen JM. School-based mindfulness instruction for urban male youth: a small randomized controlled trial. *Prev Med*. 2013;57(6):799–801
21. Sibinga EM, Webb L, Ghazarian SR, Ellen JM. School-based mindfulness instruction: an RCT. *Pediatrics*. 2016;137(1):e20152532
22. Barnes VA, Kapuku GK, Treiber FA. Impact of transcendental meditation on left ventricular mass in African American adolescents. *Evid Based Complement Alternat Med*. 2012;2012:923153
23. Sibinga EMS, Kemper KJ. Complementary, holistic, and integrative medicine: meditation practices for pediatric health. *Pediatr Rev*. 2010;31(12):e91–e103
24. Flook L, Smalley SL, Kitil MJ, et al. Effects of mindful awareness practices on executive functions in elementary school children. *J Appl Sch Psychol*. 2010;26(1):70–95
25. Barnes VA, Treiber FA, Johnson MH. Impact of transcendental meditation on ambulatory blood pressure in African-American adolescents. *Am J Hypertens*. 2004;17(4):366–369
26. Barnes VA, Bauza LB, Treiber FA. Impact of stress reduction on negative school behavior in adolescents. *Health Qual Life Outcomes*. 2003;1(10):1–7
27. Hagins M, Haden SC, Daly LA. A randomized controlled trial on the effects of yoga on stress reactivity in 6th grade students. *Evid Based Complement Alternat Med*. 2013;2013:607134
28. Telles S, Singh N, Bhardwaj AK, Kumar A, Balkrishna A. Effect of yoga or physical exercise on physical, cognitive and emotional measures in children: a randomized controlled trial. *Child Adolesc Psychiatry Ment Health*. 2013;7(1):1–16
29. Khalsa SB, Hickey-Schultz L, Cohen D, Steiner N, Cope S. Evaluation of the mental health benefits of yoga in a secondary school: a preliminary randomized controlled trial. *J Behav Health Serv Res*. 2012;39(1):80–90
30. Nidhi R, Padmalatha V, Nagarathna R, Amritanshu R. Effect of holistic yoga program on anxiety symptoms in adolescent girls with polycystic ovarian syndrome: a randomized control trial. *Int J Yoga*. 2012;5(2):112–117
31. White LS. Reducing stress in school-age girls through mindful yoga. *J Pediatr Health Care*. 2012;26(1):45–56
32. Mendelson T, Greenberg MT, Dariotis JK, Gould LF, Rhoades BL, Leaf PJ. Feasibility and preliminary outcomes of a school-based mindfulness intervention for urban youth. *J Abnorm Child Psychol*. 2010;38(7):985–994
33. Green E, Green A, Walters ED. Voluntary control of internal states: psychological and physiological. *J Transpers Psychol*. 1970;2(1):1–26
34. Culbert T. Biofeedback with children and adolescents. In: Schaefer C, ed. *Innovative Psychotherapy in Child and Adolescent Therapy*. 2nd ed. Hoboken, NJ: Wiley; 1999
35. Schwartz MS, Andrasik FE, eds. *Biofeedback: A Practitioners Guide*. 4th ed. New York, NY: Guilford Press; 2015
36. Attanasio V, Andrasik F, Burke C. Clinical issues in utilizing biofeedback with children. *Clin Biofeedback Health*. 1985;8(2):134–141

37. Biofeedback Certification International Alliance. Become board certified. Available at: www.bcia.org/i4a/pages/index.cfm?pageid=1. Accessed December 10, 2015
38. Liu C, Zhu Q, Holroyd KA, Seng EK. Status and trends of mobile-health applications for iOS devices: a developer's perspective. *J Syst Softw*. 2011;84(11):2022–2033
39. Andrasik F, Schwartz MS. Behavioral assessment and treatment of pediatric headache. *Behav Modif*. 2006;30(1):93–113
40. Nestoriuc Y, Martin A, Rief W, Andrasik F. Biofeedback treatment for headache disorders: a comprehensive efficacy review. *Appl Psychophysiol Biofeedback*. 2008;33(3):125–140
41. Culbert TP, Banez GA. Integrative approaches to childhood constipation and encopresis. *Pediatr Clin North Am*. 2007;54(6):927–947, xi
42. Culbert TP, Banez GA. Wetting the bed: integrative approaches to nocturnal enuresis. *Explore (NY)*. 2008;4(3):215–220
43. Palsson OS, Heymen S, Whitehead WE. Biofeedback treatment for functional anorectal disorders: a comprehensive efficacy review. *Appl Psychophysiol Biofeedback*. 2004;29(3):153–174
44. Weydert JA, Ball TM, Davis MF. Systematic review of treatments for recurrent abdominal pain. *Pediatrics*. 2003;111(1):e1–e11
45. Bolek JE. Use of multiple-site performance-contingent SEMG reward programming in pediatric rehabilitation: a retrospective review. *Appl Psychophysiol Biofeedback*. 2006;31(3):263–272
46. Brüttsch K, Koenig A, Zimmerli L, et al. Virtual reality for enhancement of robot-assisted gait training in children with central gait disorders. *J Rehabil Med*. 2011;43(6):493–499
47. Wang M, Reid D. Virtual reality in pediatric neurorehabilitation: attention deficit hyperactivity disorder, autism and cerebral palsy. *Neuroepidemiology*. 2011;36(1):2–18
48. Vernon D, Frick A, Gruzeliar J. Neurofeedback as a treatment for ADHD: a methodological review with implications for future research. *J Neurother*. 2004;8(2):53–82
49. Lehrer P, Feldman J, Giardino N, Song HS, Schmalig K. Psychological aspects of asthma. *J Consult Clin Psychol*. 2002;70(3):691–711
50. Carter JL, Russell HL. Use of EMG biofeedback procedures with learning disabled children in a clinical and an educational setting. *J Learn Disabil*. 1985;18(4):213–216
51. Hoy MM, Egan MY, Feder KP. A systematic review of interventions to improve handwriting. *Can J Occup Ther*. 2011;78(1):13–25
52. Barowsky EI, Moskowitz J, Zweig JB. Biofeedback for disorders of initiating and maintaining sleep. *Ann N Y Acad Sci*. 1990;602(1):97–103
53. Morin CM, Bootzin RR, Buysse DJ, Edinger JD, Espie CA, Lichstein KL. Psychological and behavioral treatment of insomnia: update of the recent evidence (1998-2004). *Sleep*. 2006;29(11):1398–1414
54. Chiarioni G, Whitehead WE. The role of biofeedback in the treatment of gastrointestinal disorders. *Nat Clin Pract Gastroenterol Hepatol*. 2008;5(7):371–382
55. Kohen DP, Olness K. *Hypnosis and Hypnotherapy With Children*. 4th ed. New York, NY: Routledge; 2011
56. Kohen DP, Kaiser P. Clinical hypnosis with children and adolescents—What? Why? How? Origins, applications, and efficacy. *Children*. 2014;1(2):74–98
57. Kohen DP, Zajac R. Self-hypnosis training for headaches in children and adolescents. *J Pediatr*. 2007;150(6):635–639
58. Sugarman LI, Wester WC. *Therapeutic Hypnosis With Children and Adolescents*. 2nd ed. Garmarthen, United Kingdom: Crown House Publishing; 2013
59. Raz A. Does neuroimaging of suggestion elucidate hypnotic trance? *Int J Clin Exp Hypn*. 2011;59(3):363–377
60. Vlieger AM, Rutten JM, Govers AM, Frankenhuys C, Benninga MA. Long-term follow-up of gut-directed hypnotherapy vs. standard care in children with functional abdominal pain or irritable bowel syndrome. *Am J Gastroenterol*. 2012;107(4):627–631
61. Astin JA, Shapiro SL, Eisenberg DM, Forsys KL. Mind-body medicine: state of the science, implications for practice. *J Am Board Fam Pract*. 2003;16(2):131–147
62. Walker LG, Walker MB, Ogston K, et al. Psychological, clinical and pathological effects of relaxation training and guided imagery during primary chemotherapy. *Br J Cancer*. 1999;80(1–2):262–268
63. Eremin O, Walker MB, Simpson E, et al. Immuno-modulatory effects of relaxation training and guided imagery in women with locally advanced breast cancer undergoing multimodality therapy: a randomised controlled trial. *Breast*. 2009;18(1):17–25
64. Kapoor VG, Bray MA, Kehle TJ. Asthma and anxiety disorders: relaxation and guided imagery as a school-based treatment. *Can J Sch Psychol*. 2007;25(4):311–327
65. Dobson CE, Byrne MW. Original research: using guided imagery to manage pain in young children with sickle cell disease. *Am J Nurs*. 2014;114(4):26–36, 37, 47
66. Forsner M, Norström F, Nordyke K, Ivarsson A, Lindh V. Relaxation and guided imagery used with 12-year-olds during venipuncture in a school-based screening study. *J Child Health Care*. 2014;18(3):241–252
67. Staples JK, Abdel Atti JA, Gordon JS. Mind-body skills groups for posttraumatic stress disorder and depression symptoms in Palestinian children and adolescents in Gaza. *Int J Stress Manag*. 2011;18(3):246–262
68. Shockey DP, Menzies V, Glick DF, Taylor AG, Boitnott A, Rovnyak V. Preprocedural distress in children with cancer: an intervention using biofeedback and relaxation. *J Pediatr Oncol Nurs*. 2013;30(3):129–138
69. Goyal M, Singh S, Sibinga EM, et al. Meditation programs for psychological stress and well-being: a systematic review and meta-analysis. *JAMA Intern Med*. 2014;174(3):357–368
70. Orme-Johnson DW, Barnes VA. Effects of the transcendental meditation technique on trait anxiety: a meta-analysis of randomized controlled trials. *J Altern Complement Med*. 2014;20(5):330–341

71. Ager K, Albrecht NJ, Cohen M. Mindfulness in Schools Research Project: exploring students' perspectives of mindfulness. *Psychology (Irvine)*. 2015;6(7):896–914
72. Whitaker RC, Dearth-Wesley T, Gooze RA, Becker BD, Gallagher KC, McEwen BS. Adverse childhood experiences, dispositional mindfulness, and adult health. *Prev Med*. 2014;67:147–153
73. Burke CA. Mindfulness-based approaches with children and adolescents: a preliminary review of current research in an emergent field. *J Child Fam Stud*. 2010;19(2):133–144
74. Zenner C, Herrnleben-Kurz S, Walach H. Mindfulness-based interventions in schools—a systematic review and meta-analysis. *Front Psychol*. 2014;5:1–20
75. Harnett PH, Dawe S. The contribution of mindfulness-based therapies for children and families and proposed conceptual integration. *Child Adolesc Ment Health*. 2012;17(4):195–208
76. Weare K. Evidence for the impact of mindfulness on children and young people. The Mindfulness in Schools Project in association with Mood Disorders Centre. 2012. Available at: <http://mindfulnessinschools.org/wp-content/uploads/2013/02/MiSP-Research-Summary-2012.pdf>. Accessed January 15, 2015
77. Kabat-Zinn J, Hanh TN. *Full Catastrophe Living: Using the Wisdom of Your Body and Mind to Face Stress, Pain, and Illness*. 15th ed. New York, NY: Bantam Dell; 2009
78. Wright LB, Gregoski MJ, Tingen MS, Barnes VA, Treiber FA. Impact of stress reduction interventions on hostility and ambulatory systolic blood pressure in African American adolescents. *J Black Psychol*. 2011;37(2):210–233
79. Kallapiran K, Koo S, Kirubakaran R, Hancock K. Effectiveness of mindfulness in improving mental health symptoms of children and adolescents: a meta-analysis. *Child Adolesc Ment Health*. 2015;20(4):182–194
80. Biegel GM, Brown KW, Shapiro SL, Schubert CM. Mindfulness-based stress reduction for the treatment of adolescent psychiatric outpatients: a randomized clinical trial. *J Consult Clin Psychol*. 2009;77(5):855–866
81. Krisanaprakornkit T, Ngamjarus C, Witoonchart C, Piyavhatkul N. Meditation therapies for attention-deficit/hyperactivity disorder (ADHD). *Cochrane Database Syst Rev*. 2010;6:CD006507
82. Benson H, Kornhaber A, Kornhaber C, LeChanu MN. Increases in positive psychological characteristics with a new relaxation-response curriculum in high school students. *J Res Dev Educ*. 1994;27(4):226–231
83. National Center for Complementary and Integrative Health. Yoga. Available at: <https://nccih.nih.gov/health/yoga>. Updated 2015. Accessed January 2015
84. Data Resource Center for Child and Adolescent Health. National profile of complementary and alternative medicine (CAM) use for children with emotional, mental or behavioral conditions or problems (2-17 years). Available at: www.childhealthdata.org/docs/drc/emb-profile_9-27-12.pdf. Updated 2012. Accessed October 2014
85. Tsao JC, Meldrum M, Kim SC, Jacob MC, Zeltzer LK. Treatment preferences for CAM in children with chronic pain. *Evid Based Complement Alternat Med*. 2007;4(3):367–374
86. Khattab K, Khattab AA, Ortak J, Richardt G, Bonnemeier H. Iyengar yoga increases cardiac parasympathetic nervous modulation among healthy yoga practitioners. *Evid Based Complement Alternat Med*. 2007;4(4):511–517
87. Berger DL, Silver EJ, Stein RE. Effects of yoga on inner-city children's well-being: a pilot study. *Altern Ther Health Med*. 2009;15(5):36–42
88. Noggle JJ, Steiner NJ, Minami T, Khalsa SB. Benefits of yoga for psychosocial well-being in a US high school curriculum: a preliminary randomized controlled trial. *J Dev Behav Pediatr*. 2012;33(3):193–201
89. Nidhi R, Padmalatha V, Nagarathna R, Amritanshu R. Effects of a holistic yoga program on endocrine parameters in adolescents with polycystic ovarian syndrome: a randomized controlled trial. *J Altern Complement Med*. 2013;19(2):153–160
90. Khalsa SB, Butzer B, Shorter SM, Reinhardt KM, Cope S. Yoga reduces performance anxiety in adolescent musicians. *Altern Ther Health Med*. 2013;19(2):34–45
91. Kuttner L, Chambers CT, Hardial J, Israel DM, Jacobson K, Evans K. A randomized trial of yoga for adolescents with irritable bowel syndrome. *Pain Res Manag*. 2006;11(4):217–224
92. Carei TR, Fyfe-Johnson AL, Breuner CC, Brown MA. Randomized controlled clinical trial of yoga in the treatment of eating disorders. *J Adolesc Health*. 2010;46(4):346–351
93. Seo DY, Lee S, Figueroa A, et al. Yoga training improves metabolic parameters in obese boys. *Korean J Physiol Pharmacol*. 2012;16(3):175–180
94. Nidhi R, Padmalatha V, Nagarathna R, Ram A. Effect of a yoga program on glucose metabolism and blood lipid levels in adolescent girls with polycystic ovary syndrome. *Int J Gynaecol Obstet*. 2012;118(1):37–41
95. Galantino ML, Galbavy R, Quinn L. Therapeutic effects of yoga for children: a systematic review of the literature. *Pediatr Phys Ther*. 2008;20(1):66–80
96. Birdee GS, Yeh GY, Wayne PM, Phillips RS, Davis RB, Gardiner P. Clinical applications of yoga for the pediatric population: a systematic review. *Acad Pediatr*. 2009;9(4):212–220
97. Kaley-Isley LC, Peterson J, Fischer C, Peterson E. Yoga as a complementary therapy for children and adolescents: a guide for clinicians. *Psychiatry (Edgmont)*. 2010;7(8):20–32
98. Kraag G, Zeegers MP, Kok G, Hosman C, Abu-Saad HH. School programs targeting stress management in children and adolescents: a meta-analysis. *J Sch Psychol*. 2006;44(6):449–472
99. Rosen L, French A, Sullivan G; RYT-200. Complementary, holistic, and integrative medicine: yoga. *Pediatr Rev*. 2015;36(10):468–474
100. Yoga Alliance. Credentialing. Available at: <https://www.yogaalliance.org/Credentialing>. Accessed December 10, 2015

Mind-Body Therapies in Children and Youth

SECTION ON INTEGRATIVE MEDICINE

Pediatrics 2016;138;

DOI: 10.1542/peds.2016-1896 originally published online August 22, 2016;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/138/3/e20161896>

References

This article cites 86 articles, 8 of which you can access for free at:
<http://pediatrics.aappublications.org/content/138/3/e20161896#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Current Policy

http://www.aappublications.org/cgi/collection/current_policy

Complementary & Integrative Medicine

http://www.aappublications.org/cgi/collection/complementary_-_integrative_medicine_sub

Section on Integrative Medicine

<http://www.aappublications.org/cgi/collection/section-on-integrative-medicine>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Mind-Body Therapies in Children and Youth SECTION ON INTEGRATIVE MEDICINE

Pediatrics 2016;138;

DOI: 10.1542/peds.2016-1896 originally published online August 22, 2016;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/138/3/e20161896>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

